



જલ્લા કૃપિ હવામાન એકમ (DAMU)

ગ્રામીણ કૃપિ મૌસમ સેવા (GKMS)

કૃપિ વિજ્ઞાન કેન્દ્ર

જુ.કૃ.યુ., જામનગર- ૩૬૧૦૦૬

તા. ૦૨-૧૦-૨૦૨૧ થી તા. ૦૬-૧૦-૨૦૨૧



(#Azadi ka Uatsav-Amrit Mahotsav)

જામનગર જલ્લા કૃપિ હવામાન બુલેટીન નં. ૭૮(૨૦૨૧) તારીખ: ૦૧-૧૦-૨૦૨૧

(જૂનાગઢ કૃપિ યુનીવર્સિટી અને ભારત મૌસમ વિભાગ દ્વારા સંપન્ન)

પાછલા અઠવાડીયાનું જામનગર નું હવામાન

હવામાન પરિબળો		
વરસાદ (મી.મી)	૦.૦-૧૪૨.૦	
મહત્તમ તાપમાન(સે.ગ્રે)	૨૮.૫-૩૩.૦	
લધુતમ તાપમાન(સે.ગ્રે)	૨૪.૦-૨૫.૫	
ભેજ(%)	સવાર	૮૨-૮૮
	બપોરબાદ	૬૮-૮૮
પવનની ગતિ (કિમી/કલાક)	૩.૨-૬.૮	

હવામાન આગાહી

ઉત્તરસૌરાષ્ટ્ર ખેતાબોહવાકીય વિભાગ - જામનગર જલ્લાની હવામાન આગાહી

તારીખ	દિવસ-૧ ૦૨-૧૦-૨૦૨૧	દિવસ-૨ ૦૩-૧૦-૨૦૨૧	દિવસ-૩ ૦૪-૧૦-૨૦૨૧	દિવસ-૪ ૦૫-૧૦-૨૦૨૧	દિવસ-૫ ૦૬-૧૦-૨૦૨૧
પેરામીટર					
વરસાદ (મી.મી.)	૨	૦	૦	૦	૦
મહત્તમ તાપમાન(સે.ગ્રે)	૩૨	૩૨	૩૩	૩૩	૩૪
લધુતમ તાપમાન(સે.ગ્રે)	૨૭	૨૭	૨૭	૨૭	૨૭
મહત્તમ ભેજ (%) સવાર	૮૨	૮૦	૮૧	૮૭	૮૪
લધુતમ ભેજ (%) બપોરબાદ	૬૭	૬૩	૬૦	૬૧	૫૨
પવનની ગતિ (કિમી/કલાક)	૧૮.૫	૧૩.૫	૧૩.૭	૧૧.૩	૧૨.૧
પવનની દિશા (ડિગ્રી)	દક્ષિણ	નૈऋત્ય	પશ્ચિમ	પશ્ચિમ	વાયવ્ય
વાદળની સ્થિતિ	વાદળાયું	મોટાભાગનું વાદળાયું	મોટાભાગનું વાદળાયું	આંશિક વાદળાયું	મોટાભાગનું વાદળાયું

નોંધ: આ આગાહી સમગ્ર જલ્લાની સરેરાશ હોય છે એટલે તેને કોઈ પણ એક જગ્યા માટે લાગુ કરવી નહીં.


ભારત સરકારશીના હવામાન ખાતા તરફથી મળેલ હવામાન માહિતી અનુસાર અત્રેના ગ્રામીણ કૃપિ મૌસમ સેવા વિભાગ, કૃપિ વિજ્ઞાન કેન્દ્ર, જામનગર તરફથી જણાવવામાં આવે છે કે ઉત્તર સૌરાષ્ટ્ર ખેત આબોહવાકીય વિસ્તારના

જામનગર જલ્લામાં આગામી સપ્તાહ દરમયાન હવામાન આંશિક વાદળાયું થી વાદળાયું રહેશે.

આગામી સપ્તાહમાં જામનગર જલ્લામાં વરસાદનું પ્રમાણ નહીંવત રહેશે. આ સમય ગાળામાં મહત્તમ તાપમાન દિવસ દરમ્યાન ૩૨-૩૪ ડિગ્રી સેલ્શીયસ જેટલું તેમજ લઘુતમ તાપમાન રાત્રી દરમ્યાન ૨૭ ડિગ્રી સેલ્શીયસ જેટલું રહેવાની સંભાવના છે. સવારના સમયે બેજનું પ્રમાણ ૮૪-૮૨ % રહેશે તથા બપોરપણીના બેજનું પ્રમાણ ૫૨-૬૭ % રહેવાની શક્યતા છે. પવનની દિશા મોટેભાગે દક્ષિણ-વાયવ્ય તરફથી રહેવાની અને પવનની ઝડપ ૧૧.૩-૧૮.૫ કીમી/કલાક રહેવાની શક્યતા છે.

NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 38 (17.09.2021 to 23.09.2021) over Agricultural region

Gujarat



Agricultural vigour is good over isolated pockets of the state.


સામાન્ય કૃષિ સલાહ

- આગામી અઠવાડિયામાં જામનગર જલ્લામાં વરસાદ પડવાની શક્યતાઓ નહીંવત છે.
- પરિપક્વ અવસ્થાએ વરપના સમયે મગફળી, બાજરો, તલ વગેરે જેવા પાકોના લાણણી કર્યો હાથ ધરવા જોઈએ.
- વરાપના સમયે કપાસ, દિવેલા અને તુવેર જેવા પાકોમાં પુરક ખાતર આપવું.
- જંતુનાશક દવાઓ, ખતરોનો છંટકાવ આકાશ ચોખ્યાં હોય ત્યારે જ કરવો.
- શિયાળુ પાકોના વાવેતર માટેનું પ્લાનિંગ કરવું જોઈએ, અને તેને અનુરૂપ જમીન તૈયાર કરવી જોઈએ.
- શિયાળુ પાકોનું વધારે વહેલું વાવેતર ન કરતા ૧૫ ઓક્ટોબર થી ૧૫ નવેમ્બર ની વર્ષેના સમયગાળામાં જ કરવું જોઈએ, જ્યારે શિયાળાની શરૂઆત થાય.

Planning for Rabi Crop Sowing

ધર્તી	<ul style="list-style-type: none"> • ધર્તિના વાવેતર માટે જમીન તૈયાર કરવી. • ટપક સિંચાય ની સગવડ હોય તો તેના દ્વારા જ પિયત આપવું જોઈએ. • સારી ગુણવત્તા વાળા બીજ વાવેતર માટે એકદા કરી લેવા જોઈએ. • વાવેતર સમયને અનુરૂપ જતો પસંદ કરવી જેવીકિ <ul style="list-style-type: none"> ✓ વહેલી વાવણી (૧૦ નવેમ્બર પહેલા): શ.જે. ડબલ્યુ-૪૬૩ ✓ સમયસર વાવણી (૧૫ થી ૨૫ નવેમ્બર: લોક-૧, શ. ડબલ્યુ-૩૬૬, ૪૫૧, ૪૬૬ ✓ મોડી વાવણી: (૨૬ નવેમ્બર થી ૧૦ ડિસેમ્બર): શ.ડબલ્યુ-૧૭૩, ૧૧, લોક-૧૧
ચાળા	<ul style="list-style-type: none"> • ચાળાના વાવેતર માટે જમીન તૈયાર કરવી. • ઓફટોબર મહિનાના અંતમાં વાવેતર કરવું. • વાવેતર માટે ગુજરાત ચાળા -૩ અને ગુજરાત ચાળા -૫ જેવી વધુ ઉત્પાદન આપતી જતો પસંદ કરવી.
જરૂર	<ul style="list-style-type: none"> • જરૂરના વાવેતર માટે જમીન તૈયાર કરવી. • નવેમ્બર મહિનામાં દંડીની શરૂઆત થયે વાવેતર કરવું. • વાવેતર માટે ગુજરાત જરૂર-૩ અને ગુજરાત જરૂર-૪ નામની જતોનું વાવેતર કરવું.
ધાણા	<ul style="list-style-type: none"> • ધાણા ના વાવેતર માટે જમીન તૈયાર કરવી. • નવેમ્બર મહિનામાં દંડીની શરૂઆત થયે વાવેતર કરવું. • વાવેતર માટે અને ગુજરાત ધાણા-૨ જેવી ભલામણ કરેલ જતોનું વાવેતર કરવું.
લસાળ	<ul style="list-style-type: none"> • લસાળ ના વાવેતર માટે જમીન તૈયાર કરવી. • નવેમ્બર મહિનામાં દંડીની શરૂઆત થયે વાવેતર કરવું. • વાવેતર માટે ગુજરાત લસાળ-૩, ગુજરાત લસાળ-૪, ગુજરાત જૂનાગઢ લસાળ-૫ ભલામણ કરેલ જતોનું વાવેતર કરવું.

પાક પ્રમાણે કૃષિ સલાહ

મુખ્યપાક	કૃષિસલાહ
ચોમાસું	• પરિપક્વ અવસ્થાએ વહેલું વાવેતર કરેલ તથા ઉભડી પ્રકારની મગફળી ઉપાડવાની કામગીરી કરવી.
મગફળી	
ચોમાસું	• ગુલાબી ઈયળના: ગુલાબી ઈયળના મોનીટરીંગ માટે પ ફીશેમેન ટ્રેપ દર હેકટરે મુકવા, જો ૮-૧૦ કૂદા એક ટ્રેપમાં જોવા મળે તો બ્યુવેરીયા બાસીયાના પાવડર ૧૦૦ ગ્રામ અને ૨૦ મિલી પ્રોફેનોફોસ પંપમાં મિક્સ કરીને વરાપે છંટકાવ કરવો.
કપાસ	 
દિવેલા	<ul style="list-style-type: none"> • વધુ વરસાદને કરને પાક નિષ્ઠળ ગયેલ હોય તો અર્ધ-શિયાળું પાકોના વાવેતરનું આયોજન કરવું. • વહેલું વાવેતર કરેલ પાકમાં પુરક ખાતર આપવું. • ઘોડિયા ઈયળ અથવા લીલી ઈયળ: ઘોડિયા ઈયળ અથવા લીલી ઈયળ ના નિયંત્રણ માટે ડાયક્લોરોવોસ ૭૬ ઈસી ૧૦ મિલી અથવા મિથાઈલ ઓ ઉમેટોન ૨૫ ઈસી ૧૫ મિલી ૧૫ લીટર પાણીમાં નાખીને વરાપે છંટકાવ કરવો.  
તલ	<ul style="list-style-type: none"> • વાતાવણ ચોખ્ખું થયા પછી પરિપક્વ અવસ્થાએ તલને વાઢીને ઉભડા કરવું.
બાજરો	<ul style="list-style-type: none"> • વરસાદી વાતાવણ પછી પરિપક્વ થયેલ કુંડાની લાણણી કરી સલામત સ્થળે સૂકમાં સૂર્ય તાપ નીચે સૂકવવા.
તુવેર	<ul style="list-style-type: none"> • વંધ્યત્વનો રોગ: તુવેરમાં સ્ટરીલીટી મોઝેક (વંધ્યત્વનો રોગ, કથીરી)ના નિયંત્રણ માટે પ્રોપરાઈટ ૫૭% ઈ.સી ૩૦-૪૦ મિલી/પાંપ વરાપે છંટકાવ કરવો.  
ચોમાસું	<ul style="list-style-type: none"> • પુરતી ખાતર આપીને વરાપના સમયે પાળા બાંધવા અને પિયત આપવું. • વરાપે આંતરાઝેડ અને નિંદામણ કરવું.
કુંગળી	<ul style="list-style-type: none"> • પુરક ખાતરનો જથ્થો સમયાંતરે આપવો. • કુંગળીમાં જંબલી ધાબા અને સુકારાનો રોગ ન લાગે તે માટે ૧૫ દિવસના અંતરે મેન્કોઝેબ ૪૦ ગ્રામ અને કાર્બેન્ડાઝીમ ૨૦ ગ્રામ ૧૫ લીટર પાણીના પર્મપમાં નાખીને ૧૫ દિવસના અંતરે વરાપે છંટકાવ કરવા.  
શાકભાજી	<ul style="list-style-type: none"> • વાવણીની કામગીરી જો બાકી હોય તો વરસાદ થયે પૂર્ણ કરવી.

પાકો

- રોગીએ છોડને ખેતરમાંથી ઉપાડીને જમીનમાં ઢાટી દેવા જોઈએ.
- શાકભાજુ પાકોને નિંદણ મુક્ત રાખવા જોઈએ.
- **પાન કોરીયું:** વેલાવાળા શાકભાજુ પાકમાં પાન કોરીયાનો ઉપદ્રવટાળવા માટે ૧૫ મીલી મીથાઈલ-ઓ-ટેમેટોન (૨૫૬૮૩) અથવા ૧૫ મીલી ડાયમીથોઓટ (૩૦ ઈસી) ૧૫ લીટરપાણીના પંપમાં નાખીને વરાપે છંટકાવ કરવો.



- **શ્રીષ્ણ:** મરચીના પાકમાં શ્રીષ્ણના નિયંત્રણ માટે ૩૦ મીલીપ્રોફેનોફ્લોસ ૧૫ લીટરપાણીમાં મિશ્રણ કરી છાંટવી. શ્રીષ્ણનો વધુ ઉપદ્રવ હોય તો સ્પીનોસાડ અથવા સ્પીનેટોરમ દવાઓ-૮મીલી પ્રતિ પંપ વરાપે છાંટવી.



- **ફળમાખી:** ફળમાખીના નિયંત્રણ માટેએસીટામીપ્રોડ ૨૦ ટકા ૧૫ ગ્રામ દવા ૧૫ લીટર પાણીમાં ભેળવી વરાપે છંટકાવ કરવો.
- **લીલી ઈયળ:** ટામેટાના પાકમાં લીલી ઈયળનો ઉપદ્રવ વધુ જોવા મળે છે તેના નિયંત્રણ માટે ઈન્ડોક્રાકાર્બ ૨૫ ઈસી ૩૦ મીલી અથવા સ્પીનોસાડ ૪૫ ઈસી પ મિલી પ્રતિ પંપ વરાપે છાંટવી.
- રોંગણુના પાકમાં જો ફળ અને ડોકાની ઈયળોનો ઉપદ્રવ હોય તો રીનાક્ષીપાયર ૧૮.૫ એસ.સી દવા પ મીલી ૧૫ લીટર પાણીમાં મિશ્ર કરીને વરાપે છંટકાવ કરવો.



- **પાનકથીરી:** શાકભાજુ પાકોમાં પાનકથીરીનો ઉપદ્રવ જોવા મળે તો ફેનાઝાકવિન ૧૦ ઈસી ૧૦ મીલી અથવા ડાયફેન્ન્યુરોન ૫૦ ડબલ્યુ પી ૧૫ ગ્રામ ૧૫ લીટર પાણીમાં પંપમાં નાખીને વરાપે છંટકાવ કરવો.



- **સફેદ માખી અને લીલી:** ભાડામાં સફેદ માખી અને લીલી પોપટીના નિયંત્રણ માટે લીઓળીનુંનેલ ૫૦ મી.લી. અથવા લીઓળીના મીજનું પ ટકા અંક નુંદાવણ ૫૦૦ મી.લી. અને ડાયફેનથાયુરોન ૫૦ ટકા વે.પા. ૧૬ગ્રામ દવાને ૧૦ લીટર પાણીમાં મિશ્ર કરીને વરાપે છંટકાવ કરવો.

પશુ

- યોગ્ય માત્રામાં આવશ્યક ક્ષાર / ખનિજ મિશ્રણ ચારો સાથે ભેળવીને પ્રાણીઓને આપવું આવશ્યક છે.
- પશુઓના રહેઠાળની જગ્યાની વ્યવસ્થિત સક્ષાઈ રાખવી અને માખીઓના ઉપદ્રવથી બચવા ક્રિનાઈલ છાંટવું અથવા લીમડાનાં પાનનો ધૂમાડો કરવો.
- પશુને દ થી ૮ કી.ગ્રા. સુકો અને ૧૫ થી ૨૦ કી.ગ્રા. લીલો ચારો આપવો. દુધાળા પશુઓને પ્રતિ લીટર દુધની ઊપજ હોય તેમને નિયમિત ૧ કી.ગ્રા. ફીડ + ૫૦ ગ્રામ ખનીજ તત્વોનું મિશ્રણ આપવું.
- ઈતરડીના નિયંત્રણ માટે ટેલ્ટામેથ્રીન અથવા એમીટ્રાલ ૨ મી.લી. એક લીટર પાણીમાં નાખીને છાંટવી.

District Agrometeorological Unit (DAMU)

GraminKrushi Mausam Seva (GKMS)

Krishi Vigyan kendra

Junagadh Agricultural University

Jamnagar-361006

Date:02-10-2021 to 06-10-2021

(#Azadi ka Uatsav-Amrit Mahotsav)



Jamnagar District Agromet Advisory Service bulletin No. 79(2021) Date: 01-10-2021

(Issued jointly by Junagadh Agricultural University and India Meteorological Department)

Significant weather for the preceding week

weather parameter		
Rain fall (mm)		0.0-142.0
Maximum temperature (°C)		29.5-33.0
Minimum temperature (°C)		24.0-25.5
Relative Humidity (%)	Morning	92-99
	Afternoon	69-98
Wind speed (Kmph)		3.2-6.9

Weather forecasting for Jamnagar District

Date	Day-1 02-10-2021	Day-2 03-10-2021	Day-3 04-10-2021	Day-4 05-10-2021	Day-5 06-10-2021
Parameters					
Rain fall (mm)	2	0	0	0	0
Maximum temperature (°C)	32	32	33	33	34
Minimum temperature (°C)	27	27	27	27	27
Relative Humidity-I (%)	92	90	91	87	84
Relative Humidity-II (%)	67	63	60	61	52
Wind speed (Km/h)	19.5	13.5	13.7	11.3	12.1
Wind Direction (deg.)	S	SW	W	W	NW
Cloud cover (Okta)	Cloudy	Generally Cloudy	Generally Cloudy	Partly Cloudy	Generally Cloudy

Note: Above forecast is the average situation of whole district not for one particular place.

Gramin Krishi Mausam Seva, Krishi Vigyan Kendra, Jamnagar and Indian meteorological department announce upcoming five days weather forecast of

Jamnagar district have Partly Cloudy to weather forecasted for next five days.


Rain fall: In the next week rainfall chances is less.

Temperature: Maximum and minimum temperature likely to be in range 32-34 °C and 27 °C, respectively.

Relative Humidity: The RH-I and RH-II is likely to be in the range of 84-92 % & 52-67 %, respectively.

Wind Direction and Speed: Direction and speed of wind is likely to be S-NW and 11.3-19.5 km/h, respectively.

**NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 38
(17.09.2021 to 23.09.2021) over Agricultural region**



Agricultural vigour is good over isolated pockets of the state.

General Agricultural Advisory

- In next week rainfall chances is less in Jamnagar.
- Harvest the crop like groundnut, bajara and sesame at maturity stage.
- Apply the supplementary dose of fertilizer to crop like Cotton, Castor and Red gram.
- **Spray pesticides and water-soluble fertilizers in clear weather condition.**
- Carried out planning of Rabi crop sowing. On the basis of crop, prepare soil and carried out cultivation.
- Not plan to early sowing Rabi crop, sowing is occur at 15th October to 15th November at ideal date of Rabi crop sowing, when cold night is starting.

Planning for Rabi Crop Sowing

Wheat	<ul style="list-style-type: none"> • Prepare the soil for wheat crop cultivation • If drip irrigation facility available then cultivate wheat crop in drip irrigation. • Collect the wheat crop seed on the basis of requirement. • Select the variety on the basis of date of sowing like <ul style="list-style-type: none"> • Early sowing: (before 10th November): GJW-463 • Timely Sowing: (15 to 25 November): Lok-1, GW-366, 496 • Late sowing: (26 November to 10 December): GW-173,11 and Lok-11
Chick pea	<ul style="list-style-type: none"> • Prepare the soil for Chick pea crop cultivation. • Sowing can be carried out in October month. • Select the high yielding variety like Gujarat chickpea-3 and Gujarat chickpea-5.
Cumin	<ul style="list-style-type: none"> • Prepare the soil for Chick pea crop cultivation. • Sowing can be carried out in November month when cold night starting. • Select the high yielding variety like Gujarat Cumin-3 and Gujarat Cumin -4.
Coriander	<ul style="list-style-type: none"> • Prepare the soil for Chick pea crop cultivation. • Sowing can be carried out in November month when cold night starting. • Select the recommended variety like Gujarat Coriander-2.
Garlic	<ul style="list-style-type: none"> • Prepare the soil for Chick pea crop cultivation. • Sowing can be carried out in November month. • Select the high yielding variety like Gujarat Garlic-3, Gujarat Garlic-4, Gujarat Junagadh Garlic-5.

Crop wise Advisory

crop	Advisory
Kharif Groundnut	<ul style="list-style-type: none"> • At maturity stage harvest the early sowing or bunch type groundnut.
Cotton	<ul style="list-style-type: none"> • Pink ballwarm: Put the 5 pheromone trap per hectare and monitor the trap, if 8-10 adult observe in trap then spray 100 gm Beauveria bassiana

powder and 20 ml prophenophos mix in 10 liter of water in clear weather condition.



- If crop is fail/ dry due to heavy rainfall then planning the resowing of other crop like half-Rabi crop.

Castor

- **semi looper and spodoptera:** For the control of semi looper and spodoptera spray dichlorovos 76 EC 10 ml or methyl o demeton 25 EC 15 ml mix in 15 liter water and spray alternate in clear weather condition.



- Apply supplementary dose fertilizer and prepare bund in castor crop and also apply irrigation when required.
- Do inter culturing and weeding at Vapsa condition.

Sesame

- After rainy period at maturity stage cut the crop in vaspa condition and make bunch for drying.

pearl millet

- After rainy period cut the mature ear head and sundry on safe place.

Red gram

- **Starility mosaic Virus:** Red mite infestation and spread Starility mosaic virus, for control them spray properzait 57% EC 30-40 ml/pump in clear weather condition.



- Apply supplementary dose fertilizer and prepare bund apply irrigation when required.
- Do inter culturing and weeding at Vapsa condition.

Kharif Onion

- At clear weather condition apply Supplementary dose of fertilizer.
- For the control of purple blotch and fuserium wilt disease in kharif onion make 3 alternate spray of Mencozeb 40 gm and Carbendazim 15 gm in 15 litre of water at 10 days interval in clear weather condition.



Vegetables

- Sowing operation is not carried then sowing operation carried out when sufficient rainfall received.
- Infected plants should be uprooted and buried in the soil.
- Keep vegetable crop weed free.
- **leaf Webber:**For the control of leaf Webber in fruit leafy vegetable spray 15 ml methyl-O-demeton (25 EC) Or 10 ml diamethoate (30 EC) in 15 liter of water in clear weather condition.



- **Thrips in chilli:**For the control of thrips in chilli, spray 30 ml profenophos in 15 litres of water. In case of heavy attack make spray of spinosad or spinetoram 7-8 ml per 15 liter of water in clear weather condition.



- **Fruit fly:**For the control of Fruit fly spray acetamiprid 20 % 15 gm per 15 liter water pump in clear weather condition.
- **Fruit borer:**In brinjal and tomato crops control of fruit borer, spray reenaxipayar @ 5 ml in 15 lit of water. In tomato crops control of fruit borer, spray indoxacarb 25 EC 30 ml or spinosad 45 EC 5 ml per pump mix in water and spray in clear weather condition.



- **Mite:**If mite infestation observe in vegetable then spray fenazaquin 10 EC 10 ml or diafenuron 50 WP 15 gm per 15 liter water pump and carried out alternate spray in clear weather condition.



- **Whitefly & jassids in Okra:** For control of whitefly &jassids in Okra, spray 500 ml of 5 % neem seed extract or 50 ml neem oil or acephate75 % SP@ 15 gm or imidacloprid 17.8 SL 3 ml/10 liter of water in clear weather condition.

Animals

- Spray phenyl in the animal shed to avoid flys and mosquitoes. Keep animal shed hygienic and clean to avoid flies and mosquitoes.
- Provide 6-8 kg good quality dry roughage with 15-20 kg green fodder to the animals. For milch animals, regularly follow schedule of 1 kg feed + 50 g mineral mixture per 2 liters of milk yield.
- Spray Deltamethrin or Amitraz 2 ml/litre of water for the control of tick.